

When Does the Risk of Not Testing Outweigh the Risk of COVID-19?

April 15, 2020

10:00 a.m. CST

Zoom Webinar

What's at Stake?


1. Patients with increased risk of COVID-19 complications
2. Increased risk of drug misuse or abuse due to isolation
3. Increased risk of drug misuse or abuse due to sudden unemployment
4. Clinicians are responsible to decide what is best for patients

Distance Solutions

1. Integrating telemedicine
2. At home drug testing
3. Drive thru drug testing
4. Safe clinic visits


Join the Senior Scientist of Genotox Laboratories, Dr. Nick Laude, as he discusses how the COVID-19 Pandemic has created unprecedented challenges for clinicians specializing in the treatment of patients in addiction recovery and pain management. Disease complexities, controlled substance prescribing, and toxicology monitoring all pose challenges to clinicians who are unable to treat patients within medical clinics.

U.S. COUNTY-LEVEL POVERTY & DRUG OVERDOSE DEATH RATES



Sources: U.S. Census Bureau Small Area Income and Poverty Estimates, CDC Small Area Estimates of Drug Mortality.

U.S. WEEKLY UNEMPLOYMENT RATES CLAIMS: PAST 50 YEARS



Source: US Bureau of Labor Statistics



Zoom Webinar **ID:** 489-175-197 **Password:** 130397

Webinar Link: [When Does The Risk of Not Testing Outweigh The Risk of COVID-19?](#)